

S A SOUTHERN
5 0 AG®

BRUSH KILLER



CONTAINS TRICLOPYR

This product is for outdoor residential use only.

Active Ingredient(s):


Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), as the triethylamine salt* 8.8%
Other Ingredients 91.2%
Contains 6.36% Triclopyr acid - 0.6 lb/ gallon.

Keep out of reach of children

CAUTION
PRECAUCION

Si usted no entiende la etiqueta, busque a alguien papa que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements continued inside of label.



Distributed By: **Southern Agricultural Insecticides, Inc.**

PALMETTO, FL. 34220 HENDERSONVILLE, N.C. 28793 BOONE, N.C. 28607
EPA Reg. No. 62719-226-829 J3 EPA Est. No. 829-FL-1

Net Contents Liquid: 1 Gallon (3.785 liters)

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes Moderate Eye Irritation * Harmful If Swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Wear protective clothing such as a long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material, and shoes and socks when applying this product.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a physician or Poison Control Center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Have product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read all Directions for Use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in original container and keep out of reach of children and pets.
Store in a cool, dry place. Avoid contamination of food and foodstuffs.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

General Information

SA-50 brand BRUSH KILLER is recommended for the control of unwanted woody plants, vines and broadleaf weeds around homes, cabins, buildings, trails, fences, walkways, rights-of-way, and other non-crop areas.

General Use Precautions and Restrictions

*Do not use on or around edible food or food crops.

*Avoid contact with eyes or clothing during and after application.

*Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

***Entry Restrictions:** Do not enter or allow others to enter the treated area until sprays have dried.

*Do not spray when the wind is blowing toward desirable plants.

*Do not apply this product directly to, or allow to drift to grapes, vegetable crops, flowers, ornamentals, or other desirable foliage. Applications should be made when wind is low and there is little chance of spray drift. Do not apply with nozzles that produce a fine-droplet spray.

*Measuring utensils and spray equipment should not be reused for purposes other than herbicide application.

Woody Plants and Vines Controlled Include: Alder, Arkansas Rose, Arrowwood, Ash, Aspen, Beech, Birch, Blackberry, Blackgum, Box Elder, Brazilian Pepper, California Rose, Cascara, Ceanothus, Cherry, Chinquapin, Choke Cherry, Cottonwood, Crataegus (Hawthorn), Douglas Fir, Dogwood, Elderberry, Elm, Hazel, Honeysuckle, Hornbeam, Kudzu, Locust, Madrone, Maples, Mesquite, Mimosa, Mulberry, Oaks, Persimmon, Pine, Poison Ivy, Poison Oak, Poplar, Raspberry, Salmonberry, Sassafras, Scotch Broom, Sumac, Sweetbay Magnolia, Sweetgum, Sycamore, Tanoak, Thimbleberry, Trumper Creeper, Tulip Poplar, Virginia Creeper, Western Hemlock, Willow, Winged Elm, Wild Grape, Wild Rose.

Perennial Broadleaf Weeds Controlled Include: Bindweed, Burdock, Canada Thistle, Chickory, Clover, Curly Dock, Dandelion, Field Bindweed, Field Horsetail Rush, Henbit, Lambsquarter, Oxalis, Plantain, Ragweed, Smartweed, Spurge, Tansy Ragwort, Vetch, Wild Lettuce, Wild Violet.

Application Methods

For best results, apply as a full coverage foliar spray when brush and weeds are fully leafed out and actively growing. Wilting or other symptoms will occur in 1-6 weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

Foliage Spray

Using a hose-end or tank sprayer, mix 8-16 tablespoons (4-8 ounces) in enough water to make one gallon mixed spray. For mature or hard-to-control plants (such as ash, blackgum, choke cherry, elm, maple, oak, pine or winged elm), or during drought conditions, use the highest rate of application indicated. Resprouting may occur the year following treatment when lower rates are used. The addition of a surfactant (spreader/sticker such as SA-50 brand Surfactant for Herbicides) enhances herbicidal action and improves distribution of the spray on hard-to-wet plant surfaces. Apply at the rate of one gallon mixed spray to approximately 500 square feet of area. Coverage may vary depending on the amount of foliage to be treated. Using coarse spray pattern and low pressure, apply as thorough foliage spray to wet all leaves. See **General Use Precautions and Restrictions**.

Stump Treatment

To kill stumps and prevent sprouting, apply the undiluted product with a paint brush or backpack sprayer to completely cover the freshly cut surface. Do not reuse paint brush. Wrap in several layers of newspaper and discard in trash.

Vine Control

When treating vines such as poison oak, poison ivy, etc. that grow on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat as directed for stump treatment.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Southern Agricultural Insecticides, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.
SOUTHERN AGRICULTURAL INSECTICIDES, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Southern Agricultural Insecticides, Inc. or the seller. All such risks shall be assumed by Buyer.

Limitations of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following: 1. Refund of purchase price paid by buyer or user for product bought, or 2. Replacement of amount of product used.

NOTICE: Buyer assumes all risk and liability of use, storage and /or handling of this material not in accordance with the terms of this label.
IN CASE OF EMERGENCY CALL: CHEMTREC 1-800-424-9300

If any part of this label is missing or for assistance you may contact us at
www.sales@southernag.com