

S A SOUTHERN
5 0 AG®

LAWN WEED KILLER WITH TRIMEC[®]

CONTROLS: DANDELIONS, CHICKWEED,
KNOTWEED, PLANTAIN, HENBIT, SPURGE
and many other broadleaf lawn weeds.

One Gallon covers up to 51,200 sq. ft.

For use on Bahiagrass, Bermudagrass,
Bentgrass, Buffalograss, Bluegrass,
Centipedegrass, Ryegrass, Fescue, St.
Augustinegrass and Zoysiagrass. (Do
not use on Floratam of St. Augustinegrass in Florida.)

ACTIVE INGREDIENTS:	BY WT.
Mecoprop, Dimethylamine Salt	5.30%
2,4-D, Dimethylamine Salt.....	3.05%
Dicamba, Dimethylamine Salt	1.29%
INERT INGREDIENTS	90.36%

THIS PRODUCT CONTAINS: TOTAL 100.00%
0.37 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid equiv. per gallon or 4.38%
0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 2.53%
0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.08%
Isomer Specific by AOAC Method 6.D01-5
TRIMEC[®] is a registered trademark of PBI/GORDON CORPORATION.



**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See Inside label for Additional
Precautionary Statements and First Aid.

WATERBASED PRODUCT KEEP FROM FREEZING

DISTRIBUTED BY: **Southern Agricultural Insecticides, Inc.**

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607
EPA Reg. No. 2217-570-829 J7 EPA EST NO. 829-FL-1

Net Contents Liquid: 1 Gallon (3.785 liters)



**STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN or CLOTHING: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-800-5556 for emergency treatment advice.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates and may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until spray has dried.

USE PRECAUTIONS:

- Do not apply this product through any type of irrigation system.
- Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area; be particularly careful with dripline of trees and other ornamental species.

Avoid spraying St. Augustinegrass when temperature is above 90° F and under conditions of high humidity. Under these conditions, some slight temporary discoloration of the St. Augustinegrass foliage may occur. This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass in cases of accidental overdose.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

WHAT IT IS, WHERE & HOW TO USE IT:

Lawn Weed Killer With Trimec[®] is a TRIMEC[®] turf herbicide formulation, a combination of three proven weed killers: 2,4-D, mecoprop and dicamba. Together, they control a wide range of broadleaf lawn weeds. This particular formulation is low in 2,4-D for maximum safety to 2,4-D sensitive grasses such as St. Augustinegrass, bentgrass and buffalograss. Spraying should be done when broadleaf weeds are small and tender for best results.

1. Irrigate the lawn 2 or 3 days prior to spraying. Do not spray lawns under drought conditions.
2. Do not water for 24 hours after treatment.
3. To avoid damage do not apply to St. Augustine or centipede grasses while emerging from winter dormancy. Also, damage may occur to these grasses if temperatures drop below 45°F within 72 hours of application.

HAND -OPERATED SPRAYER INSTRUCTIONS

ST.AUGUSTINEGRASS, BENTGRASS & BUFFALOGRASS: Mix 1 fl. oz. in 1 gallon water and uniformly spray 400 square feet. If necessary, repeat treatment in 2 to 3 weeks if hard-to-kill weeds persist.

Do not use on 'Floritam' St. Augustinegrass in Florida.

BERMUDAGRASS, BAHIAGRASS, KENTUCKY BLUEGRASS, CENTIPEDEGRASS, FESCUES PERENNIAL RYEGRASS & ZOYSIAGRASS:

Mix 1 1/2 fl. oz. in 1 gallon water and uniformly spray over 400 square feet. On hard-to-kill species, or on older weeds, mix 2 fl. oz. in 1 gallon water and spray on 400 square feet.

WEEDS: Bedstraw, Black Medic, Buckhorn, Bull Thistle, Burdock, Buttercup, Chicory, Chickweed, Clover, Cockleburr, Dandelion, Dock, Galinsoga, Ground Ivy, Hawkweed, Heal-All, Henbit, Knotweed, Lambsquarter, Lespedeza, Mallow, Morningglory, Mustard, Oxalis, Peppergrass, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Ragweed, Sheep Sorrel, Shepherdspurse, Speedwell, Tansy Mustard, Thistle, Spurge, Velvet Leaf, Wild Carrot, Wild Garlic, Wild Lettuce, Wild Onion, Yarrow and many other broadleaf lawn weeds.

HOSE-END SPRAYER INSTRUCTIONS: Use a hose-end sprayer designed for lawns. Pour recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

For Kentucky Bluegrass, Fescue, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass and Zoysiagrass:

Product Amt. to Use: (Tbsp)	Product Amt. to Use: (Fl.oz.)	Water Fill Bottle to: (Gallons)	Area to Treat: (Sq. Feet)
3	1.5	1	400
6	3	2	800
12	6	4	1600
18	9	6	2400

For St. Augustinegrass*, Bentgrass and Buffalograss:

Product Amt. to Use: (Tbsp)	Product Amt. to Use: (Fl.oz.)	Water Fill Bottle to: (Gallons)	Area to Treat: (Sq. Feet)
2	1	1	400
4	2	2	800
8	4	4	1600
12	6	6	2400

*Do not use on Floritam of St. Augustinegrass in Florida.

CONVERSION TABLE

1 Tablespoon (Tbs.) = 3 teaspoons (tsp.), 1 Fluid Ounce (fl. oz.) = 2 Tablespoons (Tbs.),

1 Cup = 8 Fluid Ounces (fl. oz.)

Limitations on broadcast applications to residential turfgrass:

the maximum application rate is 4.0 fl. oz. of product per 1,000 sq. ft. per application (0.30 lbs 2,4-D ae, 0.50 lb. MCP-P ae, and 0.12 lb. dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl. oz. of product per 1,000 sq. ft. (0.60 lbs. 2,4-D ae, 1.0 lb MCP-P ae and 0.25 dicamba ae per acre).

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

For information or questions contact our website at Southernag.com or email us at info@southernag.com

