

**S A SOUTHERN  
5 O AG.**

# CITRUS NUTRITIONAL SPRAY



Use  
1 Tablespoonful  
per Gallon of Spray

FOR USE ON CITRUS,  
AVOCADOS, MANGOS  
AND OTHER  
TROPICAL FRUITS.

Net Contents:  
1 Pint (473 ml)

Net Wt.: 1 lbs. 4.5 Ozs.

Southern Agricultural  
Insecticides, Inc.

P.O. Box 218 Palmetto, FL 34220 F672



#### CITRUS NUTRITIONAL SPRAY

##### GUARANTEED ANALYSIS:

Total Magnesium (Mg) .....	1.00%
1.00% Water Soluble Magnesium (Mg) .....	
Sulfur (S) .....	4.10%
4.10% Combined Sulfur (S) .....	
Iron (Fe) .....	1.20%
1.20% Water Soluble Iron (Fe) .....	
Manganese (Mn) .....	1.20%
1.20% Water Soluble Manganese (Mn) .....	
Zinc (Zn) .....	1.70%
1.70% Water Soluble Zinc (Zn) .....	

Derived from: Magnesium Sulfate, Ferrous Sulfate, Man-  
ganeose Sulfate, Zinc Sulfate.

#### CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin and eyes.  
Avoid breathing spray mist. Do not take internally.

This product WILL SEVERELY STAIN  
concrete, masonry and painted sur-  
faces. Be extremely careful not to allow spray or  
runoff to contact with these materials. Do not mix  
with other fertilizers, fungicides or insecticides unless  
compatibility is known.

For questions concerning this product, please  
email us at [info@southernag.com](mailto:info@southernag.com) or consult our  
website at [www.southernag.com](http://www.southernag.com).

Information regarding the contents and levels  
of metals in this product is available at  
<http://www.aapfco.org/metals.htm>

#### DIRECTIONS FOR USE

##### CITRUS, AVOCADOS, MANGOS, TROPICAL FRUITS & ORNAMENTALS

Citrus Nutritional Spray provides the minor elements  
most needed by Citrus and Tropical Fruits.

This product will benefit the plant at any time when  
deficiency symptoms appear. It is most effective when  
applied between December and February up to the  
time flowering is completed.

Apply spray to thoroughly cover to the point of run-off.  
(preventative rate may also be used on ornamentals)

##### For prevention of deficiencies:

Use 1 tablespoonful per gallon of water.

1-2 applications per year are normally sufficient.

##### For correction of a known deficiency:

Use 2 tablespoonfuls per gallon of water and make  
at least 2 applications at two week intervals.

On mature trees, the amount of spray required is  
approximately 1 gallon for each foot of height plus 5  
gallons. (20 foot tree = 25 gal. of spray)

**NOTE:** Deficiency symptoms in many plants can be an  
indication of improper pH. Soil pH should be checked  
and adjusted if necessary.

#### FIRST AID

**IF SWALLOWED:** Drink 2 glasses of water or milk  
and seek medical attention. **EYE OR SKIN CON-  
TACT:** Flush thoroughly with a large volume of  
water. If irritation persists seek medical attention.

Buyer assumes all risks of use, storage or handling of  
this material not in strict accordance with directions and  
precautions given herewith.