



HOME PEST CONTROL SPRAY

**WITH APPLICATOR
CONTAINS PERMETHRIN**

ACTIVE INGREDIENT:

Permethrin	0.25%
OTHER INGREDIENTS:	99.75%
TOTAL:	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. 34220 Hendersonville, N.C. 28793
Boone, N.C. 28607
E.P.A. Reg. No. 829 - 295 EPA Est. No. 829-FL-1

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.
Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Environmental Hazards

This product is extremely toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, or for commercial seed production or for research purposes.

INDOOR APPLICATION

HOUSEHOLD PESTS: For spot application only. Do not use as a space spray indoors. Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer.

SURFACE SPRAYING: For 60 day control of Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover mites, Cheese mites, Granary weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadellles: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For silverfish, spray bookcases. For Ants, spray trails, nests and points of entry. Allow all treated articles to dry thoroughly before use.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboard. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboard, moldings and floors. Repeat treatment as needed.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboard, moldings, window and door frames, and entire areas of the floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use.

LIVESTOCK PESTS

Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOODSUCKING LICE: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the Bedbugs and Mites hide.

PET USE APPLIES TO USE ONLY ON DOGS

ON DOGS; TO CONTROL FLEAS, TICKS AND LICE: Ready to use. Affix applicator to container for application. Do not transfer to another container. Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat application every two weeks if necessary.
DO NOT USE ON CATS OR KITTENS.

OUTDOOR PEST CONTROL

TO HELP CONTROL TICKS IN OUTDOOR ACTIVITY AREAS: Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer. Treat a band of vegetation and leaf litter 6 - 10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

PESTS ON THE OUTDOOR SURFACE OF BUILDINGS: To temporarily control House Flies, Mosquitoes, Cluster Flies, Gnats and Flying Moths. For outdoor use only as an aid in reducing annoyance from these insects. Dilute before using. Use clean sprayer. **Carefully measure and mix 1 fluid ounce of Home Pest Control and 12 fluid ounces of water (10 ozs. per gallon).** Using dilute mixture, spray pests on outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Repeat as necessary. Flush sprayer with clean water after each use.

ORNAMENTAL FLOWER GARDENS

Ready to use. Affix applicator to container for application. Do not transfer to another container unless it is a proper pesticide applicator/sprayer.

For use on **Roses, African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, rubber plants, snapdragons, stocks, wandering jew, zinnias and other ornamentals** to kill **Whiteflies, Aphids, Lace bugs, Leafminers, Japanese Beetles, Pavement Ants, Clover Mites, Red Mites, Spider Mites, exposed Thrips, and Armyworms.**

Use sufficient spray to obtain full coverage of all foliage, but do not spray to the point of runoff. When spraying plants, use a sweeping motion spraying no closer than 18 inches from the plant. Avoid heavy wetting of tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90° F. Contact insects directly whenever possible. Repeat treatment as infestation occur.

SHAKE WELL BEFORE USING

For information and questions concerning this product please contact us by email at support@southernag.com. For emergencies concerning this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378. Have product container or label with you when calling a poison control center or doctor or going for treatment.