



SNAIL & SLUG BAIT



CONTROLLED RELEASE PELLET TO CONTROL SLUGS AND SNAILS IN TREES, VEGETABLES, SHRUBS, FLOWERS AND ORNAMENTALS

ACTIVE INGREDIENT:

Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxocyclo-octane) ..	3.25%
OTHER INGREDIENTS	96.75%
TOTAL	100.00%



KEEP OUT OF REACH OF CHILDREN WARNING

This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area when bait is present. Read and follow all directions.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. Avoid breathing dust. Avoid contact with eyes, skin, and clothing. This pesticide may be fatal to dogs or other pets if eaten. Dogs may be attracted to this bait. Confine pets during application to prevent them from believing they are being fed. Keep pets out of treated area when bait is present.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Follow directions and use with care.

This product is sold by weight not volume. Some settling of contents may have occurred during shipping and handling. EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No.

Distributed by:

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607
EPA REG. NO 71096-7-829 J1

Net Weight: 1 lb. (.45 Kg)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. READ ENTIRE LABEL. USE IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

Snail & Slug Bait is a soil surface treatment for control of snails and slugs in citrus groves, orchards, berry crops, vegetable crops and gardens, ornamental gardens, seed grasses, dichondra and turf grass lawns and in ornamental greenhouses. When applied as directed, Snail & Slug Bait will withstand rain and sprinkling and will offer protection to gardens and lawns for several weeks.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof material Category A (e.g. barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyvinylchloride (PVC or vitron), and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

AGRICULTURAL USE REQUIREMENTS


Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: long-sleeved shirt and long pants, chemical resistant gloves made of any water proof material Category A (e.g. barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC or vitron), and shoes plus socks.

NON-AGRICULTURAL REQUIREMENTS

The requirements in this box apply to uses of the product that are NOT within the scope of the Worker Protection Standard, for agricultural pesticides 40 CFR Part 70. The WPS applies only when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Confine pets during application to prevent them from believing they are being fed. Keep pets out of treated area when bait is present.

USE PRECAUTIONS

Read entire label before using. Use only as a baiting treatment to be applied directly to the soil area beneath plants as listed on this label. Do not allow bait pellet to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops. **NOTE:** Snail & Slug Bait should NOT be applied to dry soil. Applications should be made following irrigation for best results. Apply any time of the day, but evening applications are preferred. Do not re-water for 48 hours after application.



0 51538 10901 7

APPLICATION INFORMATION: For control of Snails and Slugs in the following areas, use 1 lb. to treat 1,100 to 1,500 sq. ft. of soil surface (equivalent to 30 to 40 lbs. pellets per acre). For smaller areas, use one level teaspoon per square yard. Apply when first plant damage appears or is anticipated and repeat as needed for control.

FRUIT TREE CROPS SUCH AS: Apples, Apricots, Avocados, Cherries, Peaches, Pears, Citrus (Oranges, Lemons, Grapefruit, etc.) Broadcast liberally under trees, around irrigation pipes, farm buildings and any other place where snails and slugs may congregate.

BERRY CROPS SUCH AS: Blackberries, Boysenberries, Loganberries, Raspberries, Strawberries. Apply only to soil surface.

VEGETABLE CROPS AND GARDENS SUCH AS: Artichoke, Asparagus, Beans, Beets, Black-eyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Carrots, Cauliflower, Celery, Corn, Cucumbers, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Lettuce, Melons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sugar Beets, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelons. Apply only to soil surface.

ORNAMENTAL PLANTS: ORNAMENTAL GARDENS, PLANTED BORDERS AND GROUND COVERS SUCH AS: Azaleas, Begonias, Camellias, Chrysanthemums, Daffodils, Daisies, Geraniums, Honeysuckle, Hydrangeas, Ice Plant, Impatiens, Iris, Marigolds, Pansies, Petunias, Rhododendron, Rose, Sweet Peas, Snapdragons, Spathiphyllum, and Tulips. Water the infested area thoroughly before application. Broadcast or sprinkle this product evenly over the infested areas and along edges of flower beds adjacent to border or ground cover plants. Also apply on ground at base of fences and walls in shady, protected locations. This product may be injurious to Daylilies and Clematis. Do not use on these plants.

DICHONDRA AND/OR TURFGRASS LAWNS: Broadcast this product directly and evenly.

SEED GRASSES: Broadcast this product directly and evenly.

ORNAMENTAL GREENHOUSES: Broadcast this product liberally around greenhouse, under benches in greenhouse and around plants; any place where snails and slugs might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant to Snail & Slug Bait however some slight tip burning has been noticed on extremely tender orchids. This product may be injurious to Daylilies and Clematis. Do not use on these plants. **NOTE:** Snail & Slug Bait has proven to have a reasonable residual control, however for heavily infested greenhouses, a second application is recommended 7 to 10 days following the first application to bring the greenhouse under control.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. Keep container tightly closed when not in use. Do not store in direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: Read these entire Directions and Conditions of Sale before using Snail & Slug Bait. **Disclaimer:** Southern Agricultural Ins., Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purpose referred to in the directions for use, subject to the inherent risk referred to above. Southern Agricultural Ins., Inc. makes no other express or implied warranty. In no case shall Southern Agricultural Ins., Inc. be liable for consequential, special, or indirect damage resulting from the use of or handling of this product.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Lot. No.