

HI IRON GRANULES PLUS

GRANULAR DRY NUTRITIONAL for the correction and prevention of iron and certain other minor element deficiencies on shrubs, trees, flowers and turfgrass.

MINIMUM GUARANTEED ANALYSIS:

Soluble Iron as Fe	16.00%
Soluble Manganese as Mn	3.00%
Soluble Zinc as Zn	2.00%
Soluble Copper as Cu	0.50%
Boron as B	0.10%
Combined Sulfur as S	15.00%
Chlorine not more than	0.10%

DERIVED FROM: Copper Sulfate, Iron Sulfate, Manganese Sulfate, Sodium Borate, and Zinc Sulfate

GUARANTEED BY: Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220

KEEP OUT OF REACH OF CHILDREN

CAUTION:

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes and skin. Wash thoroughly after using.

DIRECTIONS FOR USE

Apply evenly by hand with gloves, with fertilizer spreader, or incorporated into a soil mix. Avoid direct contact with foliage since injury may result. **KEEP OFF SIDEWALKS, PORCHES AND ALL MASONRY AND PAINTED SURFACES OR SERIOUS STAINING WILL RESULT.**

LANDSCAPE USES

SHRUBS AND GROUND COVER: Apply evenly in bed areas or under shrubs.

For prevention of deficiencies: Use 1/4 cup per 100 sq. ft. (5' x 20')

For correction of deficiencies: Use 1/2 cup per 100 sq. ft.

TREES: Apply 1/4 cup per inch of trunk diameter within dripline.

Wash off foliage and water in thoroughly. Persistent problems with iron chlorosis are usually associated with high soil pH. Have soil tested and treat with **Soil**

Acidifier (Sulfur) in conjunction with this product. If rapid greening is desired,

Chelated Liquid Iron or **Turf Green** may also be used.

TURFGRASS APPLICATIONS

Use 2-3 pounds per 1000 sq. ft. (20' x 50') for prevention of deficiencies and greener turfgrass. For severely yellowed turfgrass, use 5 pounds per 1,000 sq. ft. Be sure foliage is dry before application, then wash off or water in as soon as possible to avoid injury.

Sweep material off of all driveways, sidewalks, and masonry surfaces before applying water. Repeat application when needed.

COMMERCIAL NURSERY APPLICATIONS

SOIL INCORPORATION: Incorporate into soil at the rate of 1 pound per cu. yard. for most crops. The liner to be transplanted should be 2" or greater. When transplanting crops that are prone to iron deficiency with a root ball equal to one quart in volume or greater, a rate of 2 pounds per cubic yard may be used.

TOPDRESS CONTAINER RATES

Container Size	Rate
1 gallon	1 teaspoon
3 gallon	2 teaspoons
7 gallon	1 tablespoon
20 gallon	1/8 cup

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28793
Boone, NC 28607

Revised 1998