
FRONT CONTAINER LABEL

S A SOUTHERN 5 0 AG_®

CITRUS **NUTRITIONAL SPRAY**



FOR USE ON CITRUS. AVOCADOS, MANGOS, AND OTHER TROPICAL FRUITS.

Use 1 Tablespoonful per Gallon of Spray

Net Contents: 1 Pint (473 ml)

Net Wt.: 1 lbs. 4.5 Ozs. (0.567 kg.)

CONTAINER **SIDE**

F672

Agricultural Insecticides, Inc. Box 218 Palmetto, FL 34220

BACK CONTAINER LABEL

 CITRUS NUTRITIONAL SPRAY

 GUARANTEED ANALYSIS:
 1.00%

 Total Magnesium (Mg)
 1.00%

 1.00% Water Soluble Magnesium (Mg)
 4.10%

 Sulfur (S)
 4.10%

Derived from: Magnesium Sulfate, Magnesium Lignin Sulfate, Ferrous Sulfate, Ferrous Lignin Sulfate , Manganese Sulfate, Manganese Lignin Sulfate, Zinc Sulfate, Zinc Lignin Sulfate.



May cause skin irritation May cause serious eye irritation

Avoid contact with skin and eyes. Avoid breathing spray mist. Do not take internally.

FIRST AID

Flush thoroughly with a large volume of water. If irritation persists, seek medical attention.

This product WILL SEVERELY STAIN concrete. masonry, and painted surfaces. Be extremely careful not to allow spray or runoff to contact with these materials. Do not mix with other fertilizers, fungicides, or insecticides unless compatibility is known.

For questions concerning this product, please email us at info@southernag.com our consult our website at www.southernag.com.

DIRECTIONS FOR USE

Citrus Nutritional Spray provides the minor elements most needed by Citrus and Tropical Fruits.

This product will benefit the plant at any time when deficiency symptoms appear. It is most effective when applied between December and February up to the time flowering is completed.

Apply spray to thoroughly cover to the point of run-off. The below preventative rate may also be used on ornamentals.

On mature trees, the amount of spray required is approximately 1 gallon for each foot of height plus 5 gallons. (20 foot tree = 25 gal. of spray)

For prevention of deficiencies:
Use 1 tablespoonful per gallon of water.
1-2 applications per year are normally sufficient.

For correction of a known deficiency:
Use 2 tablespoonfuls per gallon of water. Make at least 2 applications at two week intervals.

NOTE: Deficiency symptoms in many plants can be an indication of improper pH. Soil pH should be checked and adjusted if necessary.

Information regarding the contents and levels of metals in this product is available at http://www.aapfco.org/metals.htm

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.