

SA SOUTHERN
50 AG®

LAWN WEED KILLER WITH TRIMEC® Broadleaf Herbicide

ACTIVE INGREDIENTS:

Mecoprop-p, Dimethylamine Salt .. 5.30%

2,4-D, Dimethylamine Salt 3.05%

Dicamba, Dimethylamine Salt 1.29%

OTHER INGREDIENTS 90.36%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

0.37 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid
equivalent per gallon or 4.38%

0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per
gallon or 2.53%

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon
or 1.08%

Isomer Specific by AOAC Method 6.DO1-5

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**

See attached booklet for complete Precautionary Statements and
Directions for Use, including First Aid and Storage and Disposal.

For use on Bahiagrass, Bermudagrass,
Bentgrass, Buffalograss, Bluegrass, Centipede-
grass, Ryegrass, Fescue, Zoysiagrass and St.
Augustinegrass.

(Do not use on Floratam St. Augustinegrass.)



**CONTROLS: DANDELIONS, CHICKWEED,
KNOTWEED, PLANTAIN, HENBIT, SPURGE and
many other broadleaf lawn weeds including
those shown in the attached booklet.**

One Quart covers up to 12,800 sq. ft.

**Net Contents:
1 Quart (.946 liters)**

Southern Agricultural Insecticides, Inc.
P.O. Box 218 Palmetto, FL 34220 F2
EPA Reg. No. 2217-576-829 EPA Est. No. 629-FL-1

ACTIVE INGREDIENTS:
 Mecoprop-p, Dimethylamine Salt 5.30%
 2,4-D, Dimethylamine Salt 3.05%
 Dicamba, Dimethylamine Salt 1.29%
 OTHER INGREDIENTS 90.36%
 TOTAL 100.00%
 THIS PRODUCT CONTAINS:
 0.37 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 4.38%
 0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 2.53%
 0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.08%
 Isomer Specific by AOAC Method 6.D01-5
 TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

LAWN WEED KILLER WITH TRIMEC®

BROADLEAF HERBICIDE

CONTROLS: DANDELIONS, CHICKWEED, KNOTWEED, PLANTAIN, HENBIT, SPURGE and many other broadleaf lawn weeds, including those shown on the inside label.

KEEP OUT OF REACH OF CHILDREN

CAUTION

WATER BASED

Southern Agricultural Insecticides, Inc.
 P.O. Box 218 Palmetto, FL 34220
 EPA REG. NO. 2217-570-829 B5
 EPA EST. NO. 829-FL-1

LIMITED WARRANTY AND DISCLAIMER. FOR USE ONLY AS DIRECTED

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY and DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

ENVIRONMENTAL HAZARDS
 This pesticide is toxic to fish and aquatic invertebrates and may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- * Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- * Do not allow people or pets to enter the treated area until sprays have dried.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage and disposal. **PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing. **PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Non refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If Partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

For information and questions concerning this product please contact us by email at info@southernag.com.

- * Do not apply as a fine mist because of potential for injury to desirable plants.
- * Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- * Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

WHAT IT IS:

Lawn Weed Killer with TRIMEC® Broadleaf Herbicide is a TRIMEC® turf herbicide formulation, a combination of three proven weed killers - 2,4-D, mecoprop-p and dicamba. Together, they control a wide range of listed broadleaf lawn weeds in residential lawns.

turfgrass:

The maximum application rate is 5.0 fl. oz. of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 fl. oz. of product per 1,000 sq. ft.

Lawn Weed Killer with TRIMEC® Broadleaf Herbicide kills major northern and southern broadleaf lawn weeds as listed, such as dandelion, chickweed, knotweed, plantain in lawns without harming lawn grasses when used as directed. .

Broadleaf Weeds Controlled

- | | |
|-------------------|---------------------|
| Aster, annual | Chickweed |
| Bedstraw | Chickweed, common |
| Beggarweed | Chickweed, mouseear |
| Bindweed | Chickory |
| Bittercress | Cinquefoil |
| Black Medic | Clover |
| Brass buttons | Clover, red |
| Buckhorn | Clover, white |
| Burclover | Cocklebur |
| Burdock | Cudweed |
| Burweed | Daisy, English |
| Buttercup | Dandelion |
| Buttonweed | Dandelion, false |
| Carolina geranium | Dock |
| Carpetweed | Dock, curly |
| Catsear | Fleabane, hairy |
| | Galinsoga |

Sprayer: Put 2.5 fl. oz. (5 Tbs.) into sprayer jar. Fill sprayer jar to the 5 gallon mark with water and spray to cover 1,000 sq. ft.

For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass And Zoysiagrass:

- **For Tank Sprayers:** Use 1 fl. oz. (2 Tbs.) per gallon of water to cover 200 sq. ft. of lawn.
- **For the Dial 'N Spray:** Measure 5 fl. ozs. (10 Tbs.) into sprayer jar for every 1,000 sq. ft. of lawn. Do not add water to the container jar. Set dial on 1 oz. and spray.
- **For Ortho Plant & Weed Sprayette® Hose-End Sprayer:** Measure 5 fl. ozs. (10 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 sq. ft.

Limitations on broadcast applications to residential

- Geranium
- Ground ivy
- Hawkweed
- Healall
- Henbit
- Henbit, red
- Knawel
- Knotweed
- Lambsquarters
- Lespedeza
- Mallow
- Matchweed
- Morningglory
- Mustard
- Mustard, Tansy
- Oxalis
- Pennywort, (creeping Charlie)
- Peppergrass
- Pigweed
- Piney woods bedstraw
- Plantain

- Plantain, broadleaf
- Plantain, narrowleaf
- Poison Ivy
- Poison Oak
- Prickly lettuce
- Puncturevine
- Purslane
- Pusley, Florida
- Ragweed
- Scarlet pimpernel
- Shepherd's purse
- Smartweed, Pennsylvania
- Sorrel, Sheep
- Speedwell
- Spurge
- Spurge, prostrate
- Spurge, spotted
- Thistle
- Thistle, bull
- Thistle, musk
- Velvet Leaf

Wild Carrot	Wild Lettuce	Wild Strawberry
Wild Garlic	Wild Onion	Yarrow

WHERE TO USE:

Lawn Weed Killer with TRIMEC® Broadleaf Herbicide is intended for use on most ornamental lawns including Kentucky Bluegrass, perennial ryegrass, fescues, bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass. Do not use this product on 'Floritam' St. Augustinegrass, carpetgrass, dichondra, or lawns containing desirable clovers. Do not spray roots of trees and ornamentals.

HOW TO TIPS:

* Spray when broadleaf weeds are small and actively growing for best results. * Be sure lawn is well watered. If dry, water 1 to 2 days before spraying. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90° F. * Rain or watering within 24 hours after treatment may reduce effectiveness. * For best results, use hose end sprayers or tank type (pump-up pressure) sprayers. Spray evenly over area at specified rates.

HOW MUCH TO USE:

Spot treatments:
For St. Augustinegrass, Bentgrass, Buffalograss and Centipedegrass:
 For Tank sprayers - Use 1/2 fluid ounce (1 Tablespoon) per gallon of water to cover 200 square feet of lawn. Do not use on Floritam or other improved varieties of St. Augustinegrass. Do not apply to St. Augustinegrass or centipedegrass emerging from winter dormancy. Temperatures below 45°F within 72 hours of application may cause damage.

For Bluegrass, Fescues, Perennial Ryegrass, Bahiagrass, Bermudagrass and Zoysiagrass: Use 1 fluid ounce (2 Tablespoons) per gallon of water to cover 200 square feet. Repeat treatment for hard-to-control weeds in 30 days if they regrow. Do not use a hose end sprayer to spray small areas or spot treat weeds.