



WETTABLE OR DUSTING SULFUR

ACTIVE INGREDIENT:

Sulfur 90.0%

INERT INGREDIENTS: 10.0%

TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF ON SKIN OR CLOTHING: *Take off contaminated clothing.
*Rinse skin immediately with plenty of water for 15 - 20 minutes.
*Call a poison control center or doctor for treatment advice.
IF IN EYES: *Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
*Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
* Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378/

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin and eyes. May cause allergic response.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:
1. long-sleeved shirt and long pants 2. chemical/water resistant gloves. 3. shoes plus socks. Goggles or face shield.
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from treated areas.

PHYSICAL OR CHEMICAL HAZARDS

When dusting, keep fire and sparks away from dust stream.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607

EPA Reg. No 829-163 J5 **Net Weight: 5 lbs.(2.27 kg.)**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. **PPE** required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long sleeved shirt and long pants, chemical/waterproof gloves, shoes plus socks and goggles or face shield.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries or greenhouses. Keep children and pets of treated areas until dust has settled or sprays have dried.

BEANS as an aid in the control of rust. Apply at 4-7 day intervals using 40 lbs. per acre as a dust, or spray to obtain thorough coverage using 16 lbs. per 100 gallons (1/2 cup full per gallon). Begin applications when rust first appears and continue thereafter through harvest.

ASTER, AZALEA, CALENDULA, CHRYSANTHEMUM (hardy), CRAPE-MYRTLE, DOGWOOD, DAHLIA, DELPHINIUM, LIGUSTRUM, LILAC, MAPLE, PHLOX, RHODODENDRON, ROSE, SNAPDRAGON AND ZINNIA to control powdery mildew. Apply to obtain thorough coverage as a dust or spray thoroughly using 4 lbs. per 100 gallons (2 tablespoonfuls per gallon). Make applications at 4 to 7 day intervals beginning when powdery mildew appears and continue application as long as disease threatens.

PEACHES to control brown rot. Apply to obtain thorough coverage using 5 to 6 lbs. per 100 gallons (3 tablespoonfuls per gallon). Apply at 3 day intervals during blossom period. Apply again as soon as all petals have fallen, and repeat when 75% of the shucks have come off. Repeat applications thereafter at 10 to 14 day intervals to harvest. Use lower rate when disease presence is light. Use higher rate during periods of high rainfall and heavy disease pressure. Applications may be continued through harvest period if brown rot is present or if humid or rainy weather occurs.

CITRUS as a supplemental control of rust mite. Apply to obtain thorough coverage. Using 5 lbs. per 100 gallons (3 tablespoonfuls per gallon) as a spray or dust using 1/2 to 1 1/2 lbs. per tree depending on size of trees. **NOTE:** Sulfur may cause injury to open blooms. Do not combine sulfur with oil sprays. Applications of sulfur should not be closer than 3 weeks to an oil application and this period should be extended during dry or cold weather. Applications during very hot weather (90°F or above) may cause injury. Before applying this product consult proper U.S.D.A. or State Agricultural Authority for specific application procedures and dosages. Do not apply more than 50 lbs. per acre to Beans, 125 lbs. per acre to Citrus or 170 lbs. per acre to Peaches in any one application. **PROCESSING NOTE:** When growing crops for processing, consult processor before applying sulfur.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in original container in a cool, dry, locked place out of reach of children and pets. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non refillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of bag in a sanitary landfill or by incineration or, if allowed by State and Local authorities, by burning. If burned, stay out of smoke

CONTAINER HANDLING : (For Residential/Household Uses): Non refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Notice: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with DIRECTIONS FOR USE and PRECAUTIONARY STATEMENTS stated on this label.

For information or questions contact our website at Southernag.com or email us at support@southernag.com

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No.



EPA Reg. No 829-163 Southern Agricultural Insecticides, Inc. P.O. Box 218, Palmetto, FL 34220