

S A SOUTHERN
5 0 AG®

24-8-16

| GUARANTEED ANALYSIS: | BY WT. |
|---|---------------------|
| TOTAL NITROGEN (N) | 24.00% |
| 5.00% Ammoniacal Nitrogen (N) | |
| 5.00% Nitrate Nitrogen (N) | |
| 14.00% Urea Nitrogen (N) | |
| AVAILABLE PHOSPHATE (P₂O₅) | 8.00% |
| SOLUBLE POTASH (K₂O) | 16.00% |
| Soluble Boron as B | 0.02% |
| Chlorine - | not more than 1.00% |
| Copper as Cu (Chelated) | 0.05% |
| Iron as Fe (Chelated) | 0.10% |
| Manganese as Mn (Chelated) | 0.05% |
| Molybdenum as Mo | 0.0005% |
| Zinc as Zn (Chelated) | 0.05% |

**SOLUBLE
FERTILIZER
PLUS
MINORS**

DERIVED FROM: Urea, Ammonium Sulfate, Ammonium Phosphates, Potassium Nitrate, Sodium Borate, Cupric Disodium EDTA, Ferric Sodium EDTA, Manganous Sodium EDTA, Sodium Molybdate, Zinc Sodium EDTA
EDTA - Ethylenediaminetetraacetate **ARTIFICIALLY COLORED**

GUARANTEED BY: Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220

CAUTION: KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed. Avoid breathing dust, and contact with skin, eyes and clothing. If ingested, give two glasses of water, induce vomiting and call a physician. If in eyes or on skin flush with water. If on clothing, launder before reuse. Wash thoroughly after handling.

Contains Potassium Nitrate, an oxidizer. Possible fire hazard.

USE PRECAUTIONS

This material is compatible with most insecticides and fungicides. Do not use in combination with Bordeaux, lime, highly alkaline materials, or nutritional calcium sprays. If compatibility is not known, test on small area before use.

Timing, method and rates of application, differences in plant varieties, growing conditions and other influencing factors are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material.

Southern Agricultural Insecticides, Inc.

P.O. BOX 218
Palmetto, FL 34220

P.O. BOX 429
Hendersonville, NC 28793

P.O. BOX 85
Boone, NC 28607

F 672

LOT NO.

Net Weight: 25 lbs.

J3