

CONTAINER FRONT LABEL

S A SOUTHERN
5 0 AG®

PALM

NUTRITIONAL SPRAY

FOR THE CORRECTION AND PREVENTION OF
YELLOWING AND "FRIZZLETOP" OF PALMS



Use 1-2
Tablespoonfuls
per Gallon
Makes up to
32 gallons

Net Contents:
1 Pint (473 ml)

Net Wt.: 1 lbs. 4.5 Ozs. (.576 kg)

CONTAINER SIDE

Southern Agricultural
Insecticides, Inc.
P.O. Box 218 Palmto, FL 34220
F672



CONTAINER BACK LABEL

PALM Nutritional

GUARANTEED ANALYSIS:

Magnesium (Mg)	1.50%
1.50% Water Soluble Magnesium (Mg)	
Sulfur (S)	3.50%
3.50% Combined Sulfur (S)	
Iron (Fe)	1.00%
1.00% Water Soluble Iron (Fe)	
Manganese (Mn)	2.50%
2.50% Water Soluble Manganese	

DERIVED FROM: Magnesium Sulfate,
Magnesium Lignin Sulfonate, Ferrous Sulfate, Iron
Lignin Sulfonate, Manganese Sulfate, Manganese
Lignin Sulfonate..



May cause skin irritation
May cause serious eye irritation

Avoid contact with skin and eyes.
Avoid breathing spray mist. Do not take internally.

FIRST AID

Flush eyes or skin thoroughly with a large volume
of water. If irritation persists seek medical attention.
If swallowed: Drink 2 glasses of water or milk and
seek medical attention.

This product WILL SEVERELY STAIN
concrete, masonry and painted sur-
faces. Be extremely careful not to allow spray or
runoff to contact with these materials. Do not mix
with other fertilizers, fungicides or insecticides unless
compatibility is known.

DIRECTIONS FOR USE - PALMS

(For best results make both Soil and Foliar applications)
Foliar Application: Mix this material at the rate of 1-2
tablespoonfuls per gallon of water (1.5-3 quarts per 100
gallons). Apply to thoroughly wet the fronds to the point
of runoff, also spray some of the material directly into
the heart of the palm. **(Other Ornamentals:** use at the
same rate and spray to the point of runoff.)

Soil Application: For small palms use 1 oz. of material
in at least one half gallon of water per tree.

For medium sized palms use 2-4 oz. of material in at
least one gallon of water per tree. For large palms use
4-8 fluid ounces in at least 2 gallons of water per tree.

Treat the root area, this area usually extends out as
far as the foliage.

Curative applications: apply every 14 days until cured.
Preventative maintenance: apply every 3-4 months.

NOTE: Deficiency symptoms in many plants are an
indication of improper pH. Soil pH should be checked
and adjusted if necessary.

Information regarding the contents and levels of
metals in this product is available at
<http://www.aapfco.org/metals.htm>

For questions concerning this product, please
email us at info@southernag.com or consult our
website at www.southernag.com.

Buyer assumes all risks of use, storage or handling
of this material not in strict accordance with
directions and precautions given herewith.

