

For use in combination with  
**ATRAZINE, POAST<sup>®</sup>**, and cer-  
tain other herbicides.

**Principal functioning agents:**

Paraffin base petroleum hydrocarbons  
..... 83%  
Oil soluble surfactant blend  
..... 17%

**CAUTION**

**Keep Out Of Reach Of Chil-  
dren**

Avoid contact with skin, eyes, and clothing.  
Direct contact with eyes or skin may cause severe  
irritation.

**FIRST AID**

**IF IN EYES:** Flush with water for at least 15 minutes  
and get prompt medical attention. **IF ON SKIN:**  
Wash skin thoroughly with soap and water; remove  
and wash clothing before reuse. **IF SWALLOWED:**  
Dilute by giving water to drink and call a physician.  
Do not induce vomiting. Never give anything by  
mouth to an unconscious person.

**Southern Agricultural Insecti-  
cides, Inc.**

Palmetto, FL 34220 Hendersonville, NC 28793

**SA SOUTHERN  
50 AG**

**HERBI-OIL  
83-17  
SPRAY ADJUVANT**

Boone, NC 28607

J3 **Net Content Liquid: 1 Pint (.474 liters)**

**INSTRUCTIONS**

For use only as directed by the herbicide label, with those herbicides for which an oil based spray adjuvant is recommended. Unless otherwise indicated on the herbicide label, fill spray tank to 2/3 of the desired amount of water, add any emulsifiable materials, then HERBI-OIL 83-17 SPRAY ADJUVANT, and then wettable powders. Allow a mixing interval to completely mix each material between each addition of material. Then fill tank to desired level with continuous agitation. This product may be used with Atrazine, Basagran<sup>®</sup>, Amine 2,4-D, Fusilade<sup>®</sup>, Poast<sup>®</sup> and Trimec.

For spot or small area treatments, a 1% solution of Oil Base Spray Adjuvant is suggested when in conformity with the herbicide label.

<b>Desired Spray Volume</b>	<b>Amount SA-50 HERBI-OIL to be added to obtain a 1% solution</b>
1 Gallon	4 tsp.
3 Gallons	2 oz. (4 Tbs.)
25 Gallons	16 oz. (1 Pint)

Check pesticide label for rates specific to that product.

**PRECAUTIONS**

Use only in a well ventilated area. This product, when used with pesticides, may enhance skin absorption of the pesticide. Do not use with strong oxidizing or reducing agents, chlorate herbicides, ammonium nitrate or fertilizers containing ammonium nitrate, potassium nitrate, or sodium nitrate. Before making a selective herbicide application, test spray a small area to determine phytotoxicity to the desired crop, unless prior experience has proven safety to the desired crop under similar application conditions. Use with caution during periods of high temperature and high humidity, or when desired crop is under stress from lack of moisture or other cause. Keep container closed when not adding contents to spray tank.

**NOTICE OF WARRANTY**

Southern Agricultural Insecticides, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of application, all of which are beyond the control of Southern Agricultural Insecticides, Inc. In no case shall Southern Agricultural Insecticides, Inc. be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer Southern Agricultural Insecticides, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.