S A SOUTHERN 5 D AG®

2,4-D AMINE WEED KILLER SELECTIVE BROADLEAF WEED CONTROL

Distributed by: Southern Agricultural Insecticides, Inc Palmetto, F134220 Hendersonville, NC 2873, Boone, NC 28607 F1 EPA Est. No. 329-FL-1 EPA Reg. No. 48273-4-829

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN DANGER-PELIGRO

Si usted no entiende la etiqueta, busque alguien para que se la explique en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside of label booklet for additional Precautionary Statements, First Aid, and Directions for Use.

FOR BROADLEAF WEED CONTROL IN: PASTURES AND RANGELAND, LAWNS, GOLF COURSES, CEMETERIES, PARKS, AND SIMILAR ORNAMENTAL TURF, FOR FENCES, DITCHBANKS,



00203

Net Contents Liquid: 1 Quart (.946 liters)

KEEP OUT OF REACH OF CHILDREN DANGER-PELIGRO

IF IN EYES: Hold eyelids open and rinse slowly and gently with water for 15-20 ninutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. IF SWAL-LOWED: Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN or CLOTHING: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center. IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center o doctor, or going for treatment. You may also contact 1-800-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO: Corrosive: causes irreversible eve damage. Harmful i swallowed or inhaled. Do not get in eyes, on skin or on clothing. Avoid breathng vapor or sprav mist

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate butyl rubber, nitrile rubber or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. All mixers, loaders, applicators, flaggers, and other handlers must wear: *Long-sleeved shirt and long pants * Shoes and socks, plus protective eveware * chemical resistant gloves, when applying post harvest dips or sprays o citrus, applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate *chemical resistant apron when applying post harvest dips or sprays to citrus, nixing or loading, cleaning up spills or equipment or otherwise exposed to the ncentrate Follow manufacturer's instructions for cleaning and maintaining PPE. If no other instructions for washing, use detergent and hot water. Keep and wash all PPE separately from other laundry. See engineering controls for additional requirements

ENGINEERING CONTROLS STATEMENTS

When handlers use enclosed cabs in a manner that meets the requirement listed in the Worker Protection Standard (WPS) for agricultural pesticides [4 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or odified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. 2. Immediately remove clothing/PPE if pesticide gets inside. Then thoroughly wash and put on clean clothing. 3. Immediately emove PPE after handling this product. Wash the outside of gloves before removing. As soon as possible, thoroughly wash and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as permitted by this label Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate

This chemical has properties and characteristics associated with chemicals detected in aroundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result n contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying. Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the ency responsible for pesticide regulation. Do not apply this product through

any type of irrigation system

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker susceptible woody plants without injury to most established grasses. Protection Standard, 40 CFR part 170. This Standard contains requirements A partial list of weeds controlled: Beggarticks, Bitterweed, Blueweed Texas, for the protection of agricultural workers on farms, forests, nurseries, and Broomweed, Buckwheat Wild, Burdock, Burhead, Carpetweed, Catnip, Chicory, eenhouses, and handlers of agricultural pesticides. It contains requirements Cocklebur. Coffeeweed, Cornflower, Croton, Dandelion, Docks, Dogfennel, for training, decontamination, notification, and emergency assistance. It also Fanweed, Galinsoga, Garlic (Wild), Goatsbeard, Halogeton, Hemp (Wild), contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry Jewelweed, Jimsonweed, Kochia, Lambsquarter, Mallow Venice, Marshelder, intervals. The requirements in this box only apply to uses of this product Mildvetch, Morningglory (Annual), Mustards, Nettles, Onion (Wild), Pennycress, that are covered by the Worker Protection Standard Pepperweed (Field), Pigweed, Plantains, Poorjoe, Radish (Wild), Ragweed, Do not enter or allow worker entry into treated areas during the restricted Rape (Wild), Redstem, Salsify, Shepherdspurse, Sicklepod, Smartweed, entry interval (REI) of 48 hours. Smartweed Bitter, Sowthistle (Annual), Spanishneedles, Sunflower, Sweetclo-PPE required for early entry to treated areas that is permitted under the ver, Tansymustard, Tansyragwort, Thistle Bull, Thistle Musk, Thistle Russian, Worker Protection Standard and that involves contact with anything that has Velvetleaf, Vervains, Vetch, Water Plantain, Witchweed, Wormwood, Yellow been treated, such as plants, soil, or water, is: 1. Coveralls 2. Waterproof Rocket, Yellow Starthistle, gloves, 3. Chemical-resistant gloves made of any water-proof material, 4

Shoes plus Socks, 5. Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

Fill the spray tank half full, then add the required amount of this product and conne requirements in this box apply to uses of this product that are **NOT** within tinue filling the tank with the balance of water. Keep agitator running when filling the scope of the Worker Protection Standard for agricultural pesticides (4 the tank and during spray operations. Apply this product as a water spray during CFR Part 170). The WPS applies when this product is used to produce warm weather when weeds or brush are actively growing. Treatment during agricultural plants on farms, forests, nurseries, or greenhouses. Do not allow spray drift. On cropland and along roadsides, do not exceed 20 psi pressure. people (other than applicator) or pets on treated area during application. D not enter treated area until sprav has dried. Apply enough spray volume to provide uniform coverage of weeds and brush, usually 5 to 20 gallons water per acre by ground equipment and 1 to 5 gallons

2,4-D AMINE WEED KILLER Selective Broadles

Weed Control ACTIVE INGREDIENT Dimethylamine salt of 2.4-D-Dichloro phenoxyacetic acid 46.8% OTHER INGREDIENTS: 53.2% TOTAL 100.0% *2,4-D-Dichlorophenoxyacetic Acid Equivalent 38.4% = 3.8 lbs/gal Isomer Specific by AOAC Method No. 6 DO1-5 For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

See inside of booklet for continued Precautionary Statements and Directions for use.

Distributed by: Southern Agricultural Insecticides

P.O. Box 218 Palmetto, FL 34220 EPA Reg. No. 48273-4-829 B3 EPA Est. No. 829-FL-1

WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER. OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INFEFECTIVENESS MAY RESULT. BECAUSE OF SUCH FACTORS AS A PRESENCE OR ABSENCE OF OTHER MATERIALS LISED IN COMBINATION WITH THE GOODS, OR THE MANNER OF LISE OR APPLICATION INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER. MANUFACTURER AND SELLER. AND NO WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. IN NO EVENT SHALL MANUFAC-TURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES. OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER. IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS, OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. USER. OR ITS CUSTOMERS, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, The directions for use of this product must be followed carefully. TO THE EXTENT CONor limitations on liability, do not use the product, and return it unopened to the Seller, and the SISTENT WITH APPLICABLE LAW. (1) THE GOODS DELIVERED TO YOU ARE FURNISHED purchase price will be refunded

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. PESTICIDE STORAGE: Store product in a cool, dry, locked place out of reach children. Store at temperatures above 32ºF.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate water is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA egional office for guidance

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this ontainer. Offer for recycling if available. Triple rinse container (or equivalent omptly after emptying. Triple rinse as follows: Empty the remaining contents nto application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 14 full with water and recap. Shake for 0 seconds. Pour rinsate into application equipment or a mix tank or store insate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities by burning. If burned stay out of smoke ANTY DISCLAIMER

"AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO

FOR FENCES, DITCHBANKS, ROADSIDES, AND INDUSTRIAL SITES Apply 2 to 3 pints in 100 gallons of water and spray weeds to point of runoff. Treat as soon in spring as possible, since young actively growing weeds are easier to control

Postemergence (annual and perennial weeds): Limited to 2 applications per year Maximum of 4 pints per acre per application. Minimum of 30 days between applications. Spot treatment permitted Postemergence (woody plants): Limited to 1 application per year Maximum of 8 pints per acre per year

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. SPOT TREATMENT OF WEEDS

Mix 2 to 3 tablespoonfuls per gallon of water and apply with a hand sprayer to control broadleaf weeds in small non-cropland areas. Add a suitable, approved surfactant to the spray mixture to insure better coverage of target weeds. Do not use the same sprayer for insecticides or fungicides.

GENERAL INFORMATION

This product is recommended for numerous broadleaf weeds and certain 2,4-D

HOW TO MIX

water by aircraft. Use higher gallonage to improve spray coverage. Use the lower dosages for young, succulent growth of sensitive weed species. Use the higher rate for less sensitive species and under conditions where control is more difficult. Do not mix with oil, atrazine, surfactant or other adjuvants unless specifically recommended on this label. Deep rooted perennial weeds such as Canada Thistle and Field Bindweed and many woody plants usually require repeated applications for best control. Do not use in a Greenhouses. Do not use the same equipment for applying other materials to 2,4-D susceptible crops as injury may result.

USE PRECAUTIONS

Do not apply this product or otherwise permit it to come into contact with cotton, grapes, fruit trees, vegetables, flowers or other desirable crop or ornamental plants that are 2,4-D sensitive. Do not permit spray mist to drift onto them, since even very small quantities of spray, which may not be visible, can cause severe injury during both growing and dormant periods. Use coarse sprays to minimize drift. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by using no more than 20 psi with flat fan or flooding flat fan nozzle tips; by not spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine droplet spray.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g. wind direction, wind speed. temperature, relative humidity) and method of application (e.g., ground, or airblast) cal pesticide drift requirements regarding application of 2,4-D herbicides. Where drought periods often will give poor results. Use low spray pressure to minimize can influence pesticide drift. The applicator must evaluate all factors and make states have more stringent regulations, they must be observed. Equipment: All appropriate adjustments when applying this product. Droplet Size: When ap- ground equipment must be properly maintained and calibrated using appropriate plying sprays that contain 2,4-D as the sole active ingredient. or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or

coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or carriers or surrogates. a volume mean diameter of 385 microns or greater for spinning atomizer nozzles. When applying sprays that contain 2,4-D mixed with other active ingredients that a nozzle height greater than 4 feet above the crop canopy. require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles. Wind Speed: Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, podies of water, known habitat for nontarget species, nontarget crops within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field. Temperature inversions: If applying at wind speeds less that 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions. Susceptible Plants: Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants. Other State and Local Requirements: Applicators must follow all state and lo-

Additional requirement for ground boom application: Do not apply with

SMALL GRAINS

Wheat, Barley, Oats, Rye

Post-emergence (BARLEY, WHEAT, RYE): Not underseeded with a Legume crop for annual weeds use 1 1/2 pints per acre. For perennial weeds use 2 pints per acre. For general weed control however use 2 pints per acre. Spray when grain is in full tiller stage (usually 4 to 8 inches tall) and weeds are small. Do not apply before the tiller stage nor from early boot up to the milk stage.

For Post-emergence weed control on OATS use 1/2, to 1 pint per acre. For pre-harvest treatment when grain is in the soft to hard dough stage spray to control large weeds that interfere with harvest operation. Do not graze or feed treated forage from treated fields within 2 weeks after treatment. Do not use treated straw for livestock feed.

SMALL GRAINS USE PRECAUTIONS

Preharvest: Limited to one preharvest appliation per crop cycle. Maximum of 0.5 lbs, ae/acre per application. The preharvest interval is 14 days. Post-emergence: Limited to one post-emergence application per crop cycle. Maximum of 2 1/2 pints per acre per application. Limited to 3 2/3 per acre per crop cycle.

SORGHUM (MILO)

Apply $\frac{2}{2}$ pint per acre when plants are 6 to 15 inches tall. A higher rate of $\frac{2}{2}$ to 1 pint per acre may be needed for some weeds but the chance of crop injury is likewise increased. Do not use with oil. Do not treat before plants are 5 inches tall nor during boot, flowering or early dough stages. If plants are taller than 8 inches, use drop nozzles to keep spray off the foliage

2 pints per acre per application

Use 4 pints per acre per application. Spot treatment: Use 4 pints per acre. Maximum of two applications per year.

Minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

LAWNS, GOLF COURSES, CEMETERIES, PARKS, AND SIMILAR ORNAMENTAL TURF

Apply 3 pints per acre (6 ¹/_o tsp. per 1000 sq. ft.) in sufficient water to give good coverage (usually between 3 to 5 gals. water per 1,000 sq. ft.) Increase this to 4 pints per acre (3 Tablespoons per 1000 sq. ft.) if hard to kill weeds (Spurge) are present. Do not apply to newly seeded lawns until grasses become well established. Injury may result if applied to Bentgrass, St. Augustinegrass, For Bitterweed, Broomweed, Docks, Kochia, Marshelder, and other broadleaf Carpetgrass, Centipedegrass, Dichondra, and Clovers. After weeds have died weeds: Use 3 to 4 pints per acre (6 ¹/₂ to 1 pint per acre (6 ¹/₂ to 2 ¹/₄ to 2 ¹/₄ to 2 ¹/₄ seed the area at 1 ¹/₂ times the recommended seeding rate. Keep soil moist through the first cutting.

LAWNS, GOLF COURSES, CEMETERIES, PARKS AND SIMILAR ORNA-MENTAL TURF

Postemergence:

Limited to 2 applications per year. Maximum of 3 pints per acre per application. The maximum seasonal rate is 6 pints per acre, excluding spot treatments.

GRASS SEED CROPS

2 to 4 pints per application. For difficult to control weeds and woody plants: give uniform coverage by ground application. Apply to established stands in the spring seedlings may be treated with the lower rate after the grasses have at least acre per crop cycle. 5 leaves. Perennial weed regrowth may be treated in the fall at the higher rate. GRASS SEED CROPS PRECAUTIONS

Observe the following: A 7 day pregrazing interval for dairy cattle and a

remove all dead vegetation. Prepare the seed bed by addition of top soil, peat species; if weeds are young and actively growing, 2 to 3 pints per acre (4 1/2 to emergence spray in the spring after the canes emerge and then through the tsp. per 1000 sq. ft.) may be needed for maximum control of some weeds moss or other soil amendments. Work into the topsoil. Wait 2 weeks then 6 1/2, tsp. per 1000 sq. ft.) may control same species. Deep rooted perennials layby stage in accordance with the State recommendations. Use 2 to 4 pints in but such rates are more likely to injure corn. If corn is over 8 inches tall, may require repeated treatments "fall-spring-fall" or "spring-fall-spring." In newly sufficient water to treat 1 acre. sprigged Coastal Bermudagrass apply 2 to 3 pints per acre (4 1/2 to 6 1/2 tsp. per SUGAR CANE USE PRECAUTIONS 1000 sq. ft.) either pre-emergence or post-emergence treatment PASTURES AND RANGELAND USE PRECAUTIONS

Do not cut forage for hav within 7 days of application.

as much as possible. Temporary crop injury may occur under conditions of high soil moisture and high air temperatures. If it is necessary to treat Corn (field and pop) Do not use treated crop as fodder for 7 days following crop at this time, use 2/, pint per acre. Varieties vary in tolerance to 2.4-D. Do not spray sensitive varieties. Consult with your local Extension Service personnel or University Specialist for this information.

SORGHUM (MILO) USE PRECAUTIONS

The preharvest interval (PHI) is 30 days. Do not permit meat or dairy ing application

Postemergence (acids, salts, and amines): Limited to 1 application per

CORN

Preemergence: Apply 2 to 3 pints per acre $(1 \frac{1}{2} - 3 \text{ TBLS})$. per 1000 sq. light sandy soils.

Post-emergence: After emergence of corn plants use 1/2 pint per acre (1 use drop nozzles to keep spray off the corn foliage as much as possible. Do not apply from the tasseling to the dough stage. Do not use with oil, atrazine or other adjuvants. Crop injury is more likely to occur if corn is arowing rapidly under high temperature and high soil moisture conditions. o reduce breakage of stalks from temporary brittleness caused by 2,4-D, delay cultivations for 8 to 10 days after treatment. NOTE: Hybrids vary in response to 2,4-D and some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Contact the seed company or the Agricultural

Limited to 2 applications per year. Maximum of 4 pints per acre per application. Minimum of 21 days between applications.

PASTURES AND RANGELAND

pre-slaughter interval for meat animals of 3 days. Do not use on bent grasses, alfalfa, clover or other legumes or on newly seeded pastures. Do not apply after Limited to one preplant or preemergence application per crop cycle. Maximum of heading begins or when the grass is in the boot to milk stage where grass seed production is desired

Postemergence: For susceptible annual and biennial broadleaf weeds: Use

Experiment Station weed specialists for this information CORN USE PRECAUTIONS

spring from the tiller to the early boot stage. Do not spray in boot stage. New application. The preharvest interval (PHI) is 7 days. Maximum of 6 pints per

Preplant or preemergence: Limited to one preplant or preemergence application per crop cycle. Maximum of 2 pints per acre per application. **Postemergence:** Limited to one postemergence application per crop cycle. Maximum of 1 pint animals to consume treated crop as fodder or forage for 30 days followper acre per application.

Corn (Sweet) Do not use treated crop as fodder for 7 days following application. The preharvest interval (PHI) is 45 days. Minimum of 21 days between applica- crop cycle. Maximum of 2 pints/acre per application. tions Maximum of 3 pints per acre per crop cycle. Preplant or preemergence: 2 pints per acre per application. Postemergence: Limited to one postemergence ft.) to soil anytime after planting but before com emerges. Do not use on application per crop cycle. Maximum of 1 pint per acre per application

SUGAR CANE

Do not harvest cane prior to crop maturity. Do not apply more than 8 pints per acre per crop cycle

Preemergence: Limited to one application per crop cycle. Maximum of 4 pints per acre per application. Postemergence: Limited to one application per crop cycle. Maximum of 4 pints per acre per application.

For moderately susceptible biennial and perennial broadleaf weeds: Use Use 1 to 2 pints per acre (2 1/4 to 4 1/2 tsp. per 1000 sq. ft.) in sufficient water to

Maximum of 8 pints per acre per year