SASOUTHERN SIDAG SULFATE OF POTASH

STRENGTHENS THE PLANT MAKING IT LESS SUSCEPTIBLE TO DISEASE AND ADVERSE CONDITIONS.

GUARANTEED ANALYSIS:

Soluble Potash (K₂O)......50% Derived from: Sulfate of Potash Chlorine (Cl) less than 2.00% **GUARANTEED BY:**

Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220 F672

KEEP OUT OF REACH OF CHILDREN

CAUTION

May be harmful if swallowed. Avoid contact with eyes and skin.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

DIRECTIONS FOR USE

LAWNS: If used with other fertilizers: Apply in the Spring and again in the early Fall at a rate of 1 pound per 1,000 sq. ft. This bag will cover 5,000 sq. ft.

If used as the sole fertilizer nutrient: Apply in the Spring and again in the early Fall at a rate of 4 pounds per 1,000 sq. ft.

Water in thoroughly after application.

TREES, SHRUBS AND FLOWERING PLANTS: Apply in the Spring and again in the Fall. For trees, apply evenly around the drip line at a rate of 1/2 lb. for each inch of trunk diameter. For shrubs and flowering plants, mix lightly into the soil at a rate of 1/4 cup per every 2 ft. of plant height. Water in thoroughly after application.

BEDDING PLANTS: Before planting: Apply at a rate of 1 lb. per 250 sq. ft. and thoroughly mix into soil. Plant and then water in thoroughly.

Established Plants: Apply at a rate of 1 lb. per 500 sq. ft. Be sure to wash any material off plants and then work lightly into the soil. Water in thoroughly after application.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone. NC 28607

Net Weight: 5 lbs. (2.27 kg.)

0 51538 12255

SIASOUTHERN SULFATE OF POTASH SIDAG.